

ALTERED NEUROLOGICAL/ INGESTION ADMINISTRATIVE ORDER

Initiate immediate supportive care:

O2 to maintain sat \geq 94%
Complete primary and secondary survey as indicated
Vital Signs including FSBG (temperature as indicated)
Cardiac Monitor
12-Lead ECG

Inclusion Criteria:

Drug Overdose, Diabetic emergencies, electrolyte imbalance, medication side effects,
history/possibility of dysrhythmias, Altered LOC of unknown etiology

Initiate IV NS/LR TKO

If normotensive, consider saline lock with NS flush
Push all meds slowly
If hypotensive or FSBS > 350, administer NS/LR 20ml/kg bolus
Reassess VS and lung sounds after every 500ml infused
Hold fluids for signs/symptoms of fluid overload
May repeat as needed for continued hypotension

FSBG <70

If patient is alert & intact gag reflex:

- Administer 1 Tube **Oral Glucose**, May repeat x1

If patient is unconscious or altered:

- **Dextrose 10% (D10)** 1ml/kg (max 250ml) IV
 - If after 10 minutes FSBG <70, MR dose x 1

If unable to establish IV:

- **Glucagon** 0.1mg/kg (max 1mg) IM, May repeat x1 after 10 minutes

Hypoventilation, pinpoint pupils or possible drug use:

Ensure adequate oxygenation and ventilation

Administer **Naloxone (Narcan)** 0.5-2mg IV, IM, IO, or IN, titrate to effect
May repeat as needed, every 2 minutes

Contact poison control 1-800-222-1222 as needed

Relay ingested substance to hospital, if known

Notification to include: Altered Neurological Administrative Order, chief complaint, unit number, patient age,
gender, and ETA to receiving facility.
Advise if patient is unstable.